

# A study of the effects of disorder in the 2D Hubbard model

*M. Enjalran, F. Hebert, R. Scalettar, S. Zhang, G. Batrouni  
and M. Kalos*

This article was submitted to  
UC Davis Colloquium, Davis, CA, June 5, 2000

**June 5, 2000**

**U.S. Department of Energy**

Lawrence  
Livermore  
National  
Laboratory

## DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This report has been reproduced  
directly from the best available copy.

Available to DOE and DOE contractors from the  
Office of Scientific and Technical Information  
P.O. Box 62, Oak Ridge, TN 37831  
Prices available from (423) 576-8401  
<http://apollo.osti.gov/bridge/>

Available to the public from the  
National Technical Information Service  
U.S. Department of Commerce  
5285 Port Royal Rd.,  
Springfield, VA 22161  
<http://www.ntis.gov/>

OR

Lawrence Livermore National Laboratory  
Technical Information Department's Digital Library  
<http://www.llnl.gov/tid/Library.html>

Presentation: Colloquium  
Department of Physics  
UC Davis

Date: Monday, June 5, 2000

Speaker: Matthew Enjalran

Title: A study of the effects of disorder in the 2D Hubbard model

We study the effects of disorder on long range antiferromagnetic correlations and the Mott gap in the half-filled, two dimensional, repulsive Hubbard model. We employ Hartree-Fock (HF) and Quantum Monte Carlo (QMC) techniques in our study of the bond and site disordered models. Results from mean field (HF) calculations are used to develop a qualitative picture of the physics and to guide our choice for input to the QMC methods. The basic properties of two QMC methods for correlated fermions are discussed, and the results from these different approaches are presented. This work was performed under the auspices of the U.S. Department of Energy by the University of California and Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

# A study of the effects of disorder in the 2D Hubbard model

Matthew Enjalran

Department of Physics, University of California, Davis  
Materials Research Institute,  
Lawrence Livermore National Laboratory

June 5, 2000

## Collaborators

Frédéric Hébert, Institut Non-Linéaire de Nice, Université de Nice-Sophia Antipolis, France

Richard Scalettar, Department of Physics, University of California, Davis

Shiwei Zhang, Department of Physics, College of William and Mary

George Batrouni, Institut Non-Linéaire de Nice, Université de Nice-Sophia Antipolis, France

Malvin Kalos, Lawrence Livermore National Laboratory


This work was performed under the auspices of the U.S. Department of Energy by the University of California and Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

## Introduction


- Main Goal: To study correlated Fermi systems in the presence of disorder. We will consider model systems with hopping and on-site disorder. Our focus is the destruction AFLRO and the behavior of the Mott gap in the presence of disorder.
- The single band 2D Hubbard model, it encapsulates many of the most interesting qualitative many body effects. It is also one of the simplest correlated models to work with.
  - The possibility of interacting fermions to order.
  - The appearance of insulating states in systems with partially filled bands.
  - $U < 0$ , superconductivity.
  - $U > 0$ , model for Metal-Insulator.
  - Boson Hubbard - a model for a supersolid.
  - 1D model is solved exactly by Bethe ansatz, an insulator for all  $U > 0$ , no MIT.
  - In infinite dimensions variants of the HM can be solved exactly, (Dynamical Mean Field Theory)
  - No solution for  $1 < d < \infty$ . Many approaches are employed for finite dimensions: MFT, QMC, RG, PT, ...
- We employ several techniques in our study:
  - Exact diagonalization of the non-interacting disordered model.
  - Restricted Hartree-Fock (rHF).
  - Unrestricted Hartree-Fock (uHF).
  - Determinant Quantum Monte Carlo (DQMC): R. Blankenbecler et al., PRD **24**, 2278 (1981).
  - Constrained Path Quantum Monte Carlo, (CPQMC): S. Zhang et al., PRL **74**, 3652 (1995).

# Hubbard Model

$$H = - \sum_{\langle i,j \rangle, \sigma} t_{i,j} (c_{i\sigma}^\dagger c_{j\sigma} + c_{j\sigma}^\dagger c_{i\sigma}) + U \sum_i (n_{i\uparrow} - 1/2)(n_{i\downarrow} - 1/2) + \sum_i (\epsilon_i - \mu)(n_{i\uparrow} + n_{i\downarrow})$$



- $c_{i\sigma} (c_{i\sigma}^\dagger)$ , creation (annihilation) operators for fermions of spin  $\sigma$
- $t_{i,j}$  random hopping matrix
- $U$ , on-site repulsion
- $\epsilon_i$  random site energy,  $\mu$  chemical potential ( $\mu = 0, 1/2$  filling)
- We will consider systems with uniform, uncorrelated disorder:



## Mean Field Treatment

Develop a qualitative picture of the 2D disordered phase diagram. A mean field decomposition of the Hubbard model,

$$n_{i\sigma} \rightarrow \langle n_{i\sigma} \rangle + \delta n_{i\sigma},$$

where

$$\delta n_{i\sigma} = n_{i\sigma} - \langle n_{i\sigma} \rangle.$$

produces two independent Hamiltonians

$$H = H_{\uparrow} + H_{\downarrow} - U \sum_i \langle n_{i\uparrow} \rangle \langle n_{i\downarrow} \rangle - (U/2 + \mu) \sum_{i\sigma} n_{i\sigma} + UN/4$$

$$H_{\sigma} = - \sum_{\langle i,j \rangle} t_{ij} (c_{i\sigma}^{\dagger} c_{j\sigma} + c_{j\sigma}^{\dagger} c_{i\sigma}) + \sum_i [U \langle n_{i\sigma} \rangle + \epsilon_i] n_{i\sigma}$$

Two mean field treatments:

rHF: Use the ansatz  $\langle n_{i\sigma} \rangle = (n + \sigma(-1)^i m)/2$ , solve the resulting equations by exact diagonalization.


uHF:  $\langle n_{i\sigma} \rangle$  can vary in an arbitrary manner. One solve  $H$  self-consistently.

Observables:

- Staggered magnetization:  $M_s = 1/N \sum_{i=1}^N (-1)^i m_i$ , where  $m_i$  is the magnetization at site  $i$ .
- Local magnetization:  $M_l = 1/N \sum_{i=1}^N |m_i|$ .
- Compressibility:  $\kappa = \frac{1}{N} \partial \langle n \rangle / \partial \epsilon = \frac{\beta}{N} (\langle N^2 \rangle - \langle N \rangle^2)$
- Inverse participation ratio:  $R^{-1} = 1/N \sum_{n,i} |\psi_i^n|^4$ , where the sum is over all states ( $n$ ) and sites ( $i$ ).

## rHF Results

Phase diagram for on-site disorder



A near linear relationship between  $V_c$  and  $U$ .

rHF equations:

$$H = H_{\uparrow} + H_{\downarrow} - \mu \sum_{i,\sigma} n_{i\sigma} + (Un/2 - U/2) \sum_{i,\sigma} n_{i\sigma} + UN(m^2 - n^2)/4$$

$$H_{\sigma} = - \sum_{\langle i,j \rangle} t_{i,j} (c_{i\sigma}^{\dagger} c_{j\sigma} + c_{j\sigma}^{\dagger} c_{i\sigma}) + \sum_i (\epsilon_i + \sigma(-1)^i Um/2) n_{i\sigma}$$

Solve  $H_{\uparrow}$  and  $H_{\downarrow}$  by ED.



## uHF Results

Procedure:

$$H = H_{\uparrow} + H_{\downarrow} - U \sum_{\mathbf{i}} \langle n_{\mathbf{i}\uparrow} \rangle \langle n_{\mathbf{i}\downarrow} \rangle$$

$$H_{\sigma} = - \sum_{\langle \mathbf{i}, \mathbf{j} \rangle} t_{\mathbf{ij}} (c_{\mathbf{i}\sigma}^{\dagger} c_{\mathbf{j}\sigma} + c_{\mathbf{j}\sigma}^{\dagger} c_{\mathbf{i}\sigma}) + \sum_{\mathbf{i}} [U \langle n_{\mathbf{i}\sigma} \rangle + \epsilon_{\mathbf{i}}] n_{\mathbf{i}\sigma}$$


1. Choose a disorder realization,  $t_{i,j}$  or  $\epsilon_i$ .
2. Choose an initial mean occupation,  $\langle n_{i\sigma} \rangle$ .
3. Diagonalize the two Hamiltonians,  $H_{\uparrow}$  and  $H_{\downarrow}$ .
4. Form a new mean occupation distribution,  $\langle n_{i\sigma} \rangle$ .
5. Substitute the new  $\langle n_{i\sigma} \rangle$  into the Hamiltonian.
6. Iterate steps 3-5 to self-consistency.
7. Measure observables.
8. Repeat steps 2-7 with a different initial condition. Test for convergence of physical quantities.
9. Repeat steps 1-8 for a different disorder realization.

Results:

- The same three phases are observed for the two types of disorder.
- The system always remains an insulator.

## uHF Results

Hopping Disorder ( $N = 36, 64, 100$ )




$M_l = M_s$  always, no spin glass  
 $R^{-1} \neq 0$ , always an insulator

AFMI

$$M_s \neq 0$$

$$\kappa = 0$$

On-Site Disorder ( $N = 36, 64, 100, 144$ )



AFAI

$$M_s \neq 0$$

$$\kappa \neq 0$$


PMAI

$$M_s = 0$$

$$\kappa \neq 0$$

Note: At  $U = 4$ , disorder does not destroy AFLRO.

# uHF Results



Evolution of the staggered magnetization and compressibility with on-site disorder

$12 \times 12$  lattice


Disorder destroys magnetic order for  $U = 2$  but not for  $U = 4$

## CPQMC Results

The CPQMC algorithm:

- $|\Psi^{(n+1)}\rangle = \exp^{-\Delta\tau H} |\Psi^{(n)}\rangle$
- $|\Psi^{(n)}\rangle = \sum_k c_k^{(n)} |\phi_k^{(n)}\rangle$ , Slater determinant basis.
- $|\Psi_T\rangle$ , trial wave function.
- $\langle \Psi_T | \phi_k^{(n)} \rangle \neq 0$ , constraint.
- The method ( $T = 0$ ) allows for investigations of the Hubbard model in regions where there is a sign problem. This is an issue for the case of on-site disorder.
- In our calculations,  $|\Psi_T\rangle$  is a disorder unrestricted Hartree-Fock state with  $U = 0.5$ .

### Hopping Disorder



Spin Wave Theory:  
Huse PRB **37**,2380(1988)  
 $S(\pi, \pi)/N = M^2/3 + O(L^{-1})$


Critical disorder  
 $V_c \approx 2.1$

Compare to DQMC result,  
Ulmke et al. PRB **55**,4149(1997)

$V_c \approx 1.6$

# CPQMC Results

## On-site Disorder





Critical Disorder:

$$V_c < 0.4$$

No DQMC or analytic data available for comparison.

# CPQMC Results

## Correlations



## Summary

- In the mean field limit, unrestricted Hartree–Fock, the disordered Hubbard model is an insulator. At appreciable interaction strengths, disorder does not destroy AFLRO. These results hold for hopping and on–site disorder.
- Monte Carlo simulations at  $U = 4$  show a loss of AFLRO with increased disorder.
- Preliminary results from CPQMC yield  $V_c \approx 2.1$  for the case of hopping disorder. This result is in reasonable agreement with that obtained via DQMC,  $V_c \approx 1.6$ .
- Preliminary results for on–site disorder indicate that the  $V_c \leq 0.4$ . The enhanced ability to form on–site pairs rapidly destroys AFLRO.

## Future Plans

- Need to study the effect of the trial wave function on our results; the extrapolated values for staggered magnetization are high.
- Complete work on the critical disorder at  $U = 4$ .
- Study the effect of the interaction strength  $U$  on the half–filled model.
- Map out the ground state phase diagram as a function of  $V$  and  $U$ .

